arXiv:1411.2306v4 [math.KT] 5 Aug 2016

A REPRESENTATION FOR ALGEBRAIC K-THEORY OF QUASI-COHERENT MODULES OVER AFFINE SPECTRAL SCHEMES

MARIKO OHARA

Abstract. In this paper, we study K-theory of spectral schemes by using locally free sheaves. Let us regard the K-theory as a functor K on affine spectral schemes. Then, we prove that the group completion $\Omega B^{\mathcal{G}}(B^{\mathcal{G}}GL)$ represents the sheafification of K with respect to Zariski (resp. Nisnevich) topology $\mathcal{G}$, where $B^{\mathcal{G}}GL$ is a classifying space of a colimit of affine spectral schemes GL_n.

1. Introduction

An ∞-category is a notion of categories up to coherent homotopy. The spectral algebraic geometry in terms of ∞-category, introduced by Lurie [12] [13], is a generalization of algebraic geometry.

In this paper, we study the K-theory on spectral schemes. For the K-theory of a category of projective modules of finite rank in the sense of spectral algebraic geometry, we construct an object in the ∞-category which represents the K-theory. In Section (1.1), we explain the representation of the K-theory in the classical algebraic geometry proved by Morel and Voevodsky [16] which motivates us to prove our main theorem (Theorem 1.2).

1.1. Background of this paper.

Morel and Voevodsky showed the following result of the K-theory in algebraic geometry.

Let S be an affine Noetherian scheme of finite Krull dimension. Let $\mathcal{H}(S)$ be a category defined in [16, Definition 2.1], which is arising from certain simplicial sheaves on a category of smooth schemes of finite type over S equipped with Nisnevich topology.

Theorem 1.1 ([16] Proposition 3.9)**.** *For the ordinary affine group schemes GL_n for $n \geq 0$, let $B(\coprod_n BGL_n)$ be a classifing sheaf in $\mathcal{H}(S)$. Let Σ_s be a simplicial suspension functor on $\mathcal{H}(S)$ and $R\Omega$ its right adjoint. Then, for a quasi-projective smooth scheme X over S, there is a canonical isomorphism*

$$\mathrm{Hom}_{\mathcal{H}(S)}(\Sigma_s^k(X), (R\Omega)B(\coprod_n BGL_n)) \cong \pi_k K(X),$$

Date: August 8, 2016.

2010 *Mathematics Subject Classification.* 18E99 (primary), 19D10 (secondary).

Key words and phrases. infinity category, derived algebraic geometry, K-theory.

where $K(X)$ is the K-theory of the ordinary category of locally free sheaves on X. In other words, $K(X)$ is represented by the object $(R\Omega)B(\coprod_n BGL_n))$ in the category $\mathcal{H}(S)$.

Since a simplicial presheaf is a model of an ∞-categorical sheaf, this theorem may be interpreted in terms of spectral algebraic geometry. We generalize this result to a certain ∞-category of sheaves as follows.

By an ∞-category with w^∞-cofibrations we mean a pointed ∞-category with a class of morphisms (called w^∞-cofibrations) satisfying certain conditions (see Definition 5.1 for details). Let $\mathcal{C}$ be an ∞-category with w^∞-cofibrations. We define the algebraic K-theory space of $\mathcal{C}$ by

$$K(\mathcal{C}) = \Omega|S_\bullet(\mathcal{C})|,$$

where $S_\bullet$ is the S-construction (cf. [4]) and $|-|$ is the geometric realization.

Let R be a connective $\mathbb{E}_\infty$-ring, and CAlg^{cn} the ∞-category of connective $\mathbb{E}_\infty$-rings. Let $\mathrm{Mod}_R^{\infty proj}$ be an ∞-category of projective R-modules of finite rank which we recall in Definition 2.1. It becomes an ∞-category with w^∞-cofibrations by Definition 5.3.

We denote by $\mathrm{CAlg}^{\mathcal{G}}$ an ∞-category CAlg^{cn} equipped with either the Zariski topology or the Nisnevich topology, and by $\mathrm{Spec}^{\mathcal{G}} R$ an object in the essential image of Yoneda functor $\mathrm{CAlg}^{\mathcal{G}} \to \mathcal{S}hv(\mathrm{CAlg}^{\mathcal{G}})$, where $\mathcal{S}hv(\mathrm{CAlg}^{\mathcal{G}})$ is the ∞-category of sheaves on $\mathrm{CAlg}^{\mathcal{G}}$ given in Section 2.

Let $\widehat{\mathcal{S}}$ be the ∞-category of not-necessary small spaces. We define a functor

$$K : (\mathrm{CAlg}^{\mathcal{G}})^{op} \to \widehat{\mathcal{S}} \tag{1.1}$$

which carries a spectral scheme $\mathrm{Spec}^{\mathcal{G}} R$ to the K-theory $K(\mathrm{Mod}_R^{\infty proj})$. Here, we denote by $(-)^{op}$ the opposite ∞-category.

We denote by $\widetilde{(-)}^{\mathcal{G}}$ the sheafification from the ∞-category of functors on $(\mathrm{CAlg}^{\mathcal{G}})^{op}$ to $\mathcal{S}hv(\mathrm{CAlg}^{\mathcal{G}})$ which we recall in Definition 2.4.

Let GL_n be an object in $\mathcal{S}hv(\mathrm{CAlg}^{\mathcal{G}})$. Let $B^{\mathcal{G}}GL = \coprod_{n\in\mathbb{N}} B^{\mathcal{G}}GL_n$ be the coproduct of the classifying sheaf $B^{\mathcal{G}}GL_n$ of GL_n, where $B^{\mathcal{G}}$ is a functor given by taking classifying sheaf which we recall in Section 4. Let $\Omega B^{\mathcal{G}}$ be a functor on $\mathcal{S}hv(\mathrm{CAlg}^{\mathcal{G}})$ defined in Definition 3.14. We denote by $\Omega B^{\mathcal{G}}(B^{\mathcal{G}}GL)$ the group completion on (cf. Definition 4.16).

Theorem 1.2 (cf. Theorem 6.3)**.** *Let $\mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(-,-)$ denote the mapping space of the ∞-category $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$ which we recall in Definition 4.1. There is an equivalence*

$$\mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\, \Omega B^{\mathcal{G}}(B^{\mathcal{G}}GL)) \simeq \widetilde{K}^{\mathcal{G}}(\mathrm{Mod}_R^{\infty proj}).$$

Remarks for Theorem 1.2. It is known that the sheaf $\widetilde{K}^{\mathcal{G}}$ is representable by an object in $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$ by the brown representability theorem in the sense of ∞-category [8,

Proposition 5.5.2.7]. The importance is that we give a concrete object which represents $\widetilde{K}^{\mathcal{G}}$.

Morel and Voevodsky used the cofinality theorem in Quillen's K-theory in the proof of Theorem 1.1. Since the cofinality theorem is not established in the K-theory of ∞-categories, their proof cannot be applied to our case directly. To avoid this problem, we treat an ∞-category $\mathrm{Mod}_R^{\infty proj}$ instead of the finitely generated projective modules.

1.2. **Outline of this paper.** This paper is organized as follows. In Section 2, we introduce the terminology of an ∞-category endowed with Grothendieck topology and sheaves on those ∞-category. In Section 3, we describe the group completion functor on $\mathcal{S}hv(\mathrm{CAlg}^{\mathcal{G}})$ explicitly in Proposition 3.15. In Section 4, we show the correspondence between the value of the affine group scheme GL_n on S and the automorphisms of S^n, where S is an $\mathbb{E}_\infty$-ring in Proposition 4.9. We also demonstrate that the classifying sheaf $B^{\mathcal{G}}GL_n$ is equivalent to the sheaf of projective modules of finite rank in Proposition 4.21. We also define the notion of Zariski connectedness and relate a certain mapping space of $B^{\mathcal{G}}GL_n$ is equivalent to the sheaf of projective modules of finite rank in Proposition 4.26. In Section 5, we recall the several notion and proposition with respect to K-theory. In Section 6, by using these results, we prove Theorem 6.3.

Acknowledgement. The author would like to express deeply her thanks to Professor Nobuo Tsuzuki for his valuable advice and checking this paper. The author would like to express her thanks to Professor Satoshi Mochizuki for his valuable comments on algebraic K-theory, especially the resolution theorem and the cofinality, and for reading this paper. The author also would like to express her thanks to Professor Yuki Kato for valuable comments to the author.

2. Preliminary

We fix the universe $\mathbb{U}$ such that $\mathbb{N} \in \mathbb{U}$. We define the Cat_∞ by $\mathbb{U}$-small ∞-category, which is locally $\mathbb{U}$-small.

Although there are a lot of languages of higher category theory, we use the same notation in Lurie's book [8] and papers [9], [12] and [10].

Let $\mathrm{Fun}(-, -)$ be an ∞-category of functors. For a pair of functors $f : \mathcal{C} \to \mathcal{D}$ and $g : \mathcal{D} \to \mathcal{C}$ between ∞-categories, according to [8, Proposition 5.2.2.8], we say that the functor f is a left adjoint to g (resp. g is a right adjoint to f) if there exists a unit map $u : id_{\mathcal{C}} \to g \circ f$ given in [8, Defnition 5.2.2.7].

Let CAlg be an ∞-category of $\mathbb{E}_\infty$-rings. We denote by $\mathbb{S}$ the initial object in CAlg, which is called the sphere spectrum. We say that a spectrum E is *connective* if $\pi_n E \simeq 0$

for $n < 0$. We denote by CAlg^{cn} a full ∞-subcategory of CAlg which consists of connective $\mathbb{E}_\infty$-ring.

For an $\mathbb{E}_\infty$-ring R, we also have an ∞-category Mod_R which is called the *∞-category of R-module* of Sp [9, Section 4.2]. Since the tensor product on the ∞-category of spectra is compatible with the geometric realizations [9, Corollary 4.8.2.19], Mod_R becomes the symmetric monoidal ∞-category by [9, Theorem 4.5.2.1]. We denote by $\otimes_R$ the tensor product on Mod_R.

Let R be an $\mathbb{E}_\infty$-ring and $a \in \pi_0 R$ an element. The localization of R with respect to a, which is denoted by $R[a^{-1}]$, is an $\mathbb{E}_\infty$-ring (see [13, Remark 2.9] and [9, 7.2.4]).

Definition 2.1. Let R be a connective $\mathbb{E}_\infty$-ring and M an R-module.

(i) We say that M is free of rank n if $M \simeq R^{\oplus n}$.

(ii) We say that M is finitely generated projective if it is a retract of finitely generated free R-modules.

(iii) We say that M is projective of rank n if it is finitely generated projective and we can choose elements $x_1, \cdots, x_m \in \pi_0 R$ such that they generate the unit ideal and each localization $M[x_i^{-1}]$ is a free module of rank n over $R[x_i^{-1}]$.

We denote by Mod_R^{nfree} (resp. Mod_R^{nproj}) an ∞-category of free (resp. projective) R-modules of rank n. We denote by Mod_R^{proj} the ∞-category of finitely generated projective R-modules.

Remark 2.2. Let $\mathrm{Mod}_R^{\infty proj}$ be an ∞-category of projective R-modules of finite rank. In general, $\mathrm{Mod}_R^{\infty proj}$ and Mod_R^{proj} is not equivalent.

As in [8], [9] and [2], to treat certain size of limits and colimits, we need to enlarge the universe and fix the size of universes properly.

We adopt the axiom of universes which allows us to consider that every cardinal can be strictly upper bounded by a strongly inaccessible cardinal. Then, there exists a bijection between strongly inaccessible cardinals and universes, and thus we can take a suitable enlargement of universe which we need.

By the axiom of universe, there exists an enlargement of universes $\mathbb{U} \in \mathbb{V}$ such that every $\mathbb{U}$-small object is also $\mathbb{V}$-small, so that Cat_∞ becomes a $\mathbb{V}$-small category.

In this paper, sometimes we need to treat the ∞-category CAlg as small, and to regard the maximal ∞-groupoid $(\mathrm{Mod}_R)^{\simeq}$ of Mod_R as a space. We also need to treat the size of limits and colimits. In these cases, we enlarge the universe as the following definition and proceed the arguments.

Definition 2.3. Let $\mathcal{S}$ be the ∞-category of $\mathbb{U}$-small spaces. Throughout this paper, we enlarge the unverse $\mathbb{U} \in \mathbb{V}$ such that Mod_R is a $\mathbb{V}$-small ∞-category. we use the notation $\widehat{\mathcal{S}}$ for $\mathcal{S}$ and $\widehat{\mathrm{Cat}_\infty}$ for Cat_∞ after changing the universe from $\mathbb{U}$ to $\mathbb{W}$.

2.1. **Sheaves and spectral affine schemes.** Let $\mathcal{X}$ be an ∞-category equipped with the Grothendieck topology in [8, Definition 6.2.2.1] and $\mathcal{C}$ an ∞-category which admits limits. We denote by $\mathcal{S}hv_{\mathcal{C}}(\mathcal{X})$ an ∞-category of $\mathcal{C}$-valued sheaves on $\mathcal{X}$ in [8, Definition 6.2.2.6].

Definition 2.4. Let $\widetilde{(-)} : \mathrm{Fun}(\mathcal{X}^{op}, \mathcal{S}) \to \mathcal{S}hv_{\mathcal{S}}(\mathcal{X})$ be an localization defined by [8, Definition 6.2.2.6], which is a left adjoint of the inclusion. For an object F of $\mathrm{Fun}(\mathcal{X}, \mathcal{S})$, we say that $\widetilde{F}$ is a sheafification of F.

Remark 2.5 ([8] Construction 6.2.2.9, Remark 6.2.2.12). For an ∞-category $\mathcal{C}$ equipped with a Grothendieck topology, and a presheaf F on $\mathcal{C}$, the sheafification $\widetilde{F}$ of F is given by the following formula: for any $C \in \mathcal{C}$,

$$\widetilde{F}(C) = \mathrm{colim}_{\mathcal{C}^{(0)}_{/C}} \mathrm{lim}_{C' \in \mathcal{C}^{(0)}_{/C}} F(C'),$$

where the first limit is the limit of the simplicial diagram associated to $C' \to C$ [8, Corollary 6.2.3.5] and the second colimit is taken over the collection of covering sieves on C.

Let $\mathbb{S}$ be a sphere spectrum. Here is a list of morphisms of $\mathbb{E}_\infty$-rings.

- Let $f : A \to B$ be a morphism of $\mathbb{E}_\infty$-rings. Recall that $f : A \to B$ is said to be flat if $\pi_0 B$ is flat $\pi_0 A$-module and the underlying map of commutative rings $\pi_0 A \to \pi_0 B$ induces the isomorphism $(\pi_i A) \otimes_{\pi_0 A} (\pi_0 B) \simeq \pi_i B$ for every integer i.
- Let $f : A \to B$ be a morphism of $\mathbb{E}_\infty$-rings. $f : A \to B$ is said to be étale if it is flat and the underlying map of commutative rings $\pi_0 A \to \pi_0 B$ is étale. The class of étale morphisms on $(\mathrm{CAlg}^{cn})^{op}$ satisfies the axiom of admissible morphisms by [12, Proposition 2.4.17].
- We say that a morphism $f : A \to B$ of $\mathbb{E}_\infty$-rings is faithfully flat if it is flat morphism and the underlying map of commutative rings $\pi_0 A \to \pi_0 B$ is faithfully flat. By a class of faithfully flat morphisms, Grothendieck topology is defined on CAlg^{op}, which is called flat topology [13, Definition 5.2, Proposition 5.4].

Recall the Grothendieck topology given by the admissibility on covering sieves from [12, Definition 1.2.1]. Let $\mathcal{G}^{Sp}_{Zar}(\mathbb{S})$ be an ∞-category with Grothendieck topology given as [13, Definition2.10]. We denote by CAlg^{Zar} an ∞-category $\mathrm{Ind}(\mathcal{G}^{Sp}_{Zar}(\mathbb{S})^{op})$ equipped with Zariski topology [12, Notation 2.2.6]. Moreover, this Zariski geometry is finitary [12, Remark 2.2.8] by the definition. (For the base change assumption, see [12, Remark 1.2.4]).

Let A be a connective $\mathbb{E}_\infty$-ring. Let CAlg^{et}_A be an ∞-category spanned by connective étale algebras over A. Let $\mathcal{C} \subset (\mathrm{CAlg}^{et}_A)^{op}$ be a sieve containing A. We say that $\mathcal{C}$ is a Nisnevich covering sieve on A if it contains a collection of morphisms $A \to A_a$ such that their underlying maps of commutative rings $\pi_0 A \to \pi_0 A_a$ determine a Nisnevich covering

defined in [11, Definition 1.1]. Let A be a connective $\mathbb{E}_\infty$-ring. We define the admissible morphisms in $(\mathrm{CAlg}^{cn})^{op}$ by the morphisms corresponding to étale morphisms in CAlg^{cn}, and the collection of admissible morphisms $A \to A_a$ generates a covering sieve on A if and only if it is a Nisnevich covering sieve. Then, it generates a Grothendieck topology on CAlg^{cn}. It is called the Nisnevich topology on CAlg^{cn}. We denote by CAlg^{Nis} an ∞-category CAlg^{cn} endowed with the Nisnevich topology.

Let $\mathrm{CAlg}^{\mathcal{G}}$ denote either CAlg^{Zar} or CAlg^{Nis}. Let us denote an object in the essential image of Yoneda functor $\mathrm{CAlg}^{\mathcal{G}} \to \mathcal{S}hv_{\mathcal{S}}(\mathrm{CAlg}^{\mathcal{G}})$ by $\mathrm{Spec}^{\mathcal{G}} R$, and we call the object an affine spectral scheme.

The following proposition is a special case of [13, Proposition 5.7].

Proposition 2.6. *Let us consider* CAlg^{op} *endowed with the flat topology. Then, in the case of* ∞*-topos* CAlg^{op} *with flat topology, a functor* F *is a sheaf if, it preserves finite products and for any covering* $X \to Y$ *in* CAlg*,*

$$F(Y) \to \lim_{\Delta} F(X_\bullet)$$

is an equivalence. Here, $X_\bullet \to Y$ *is a simplicial object associated to* $X \to Y$*, and the limit in the right hand side is taken over its simplicial diagram.*

Note that a Zariski (resp. Nisnevich) covering sieve is a covering which contains a finite collection of morphisms $B \to B_a$ generating the covering sieve.

Lemma 2.7. *(i) A Zariski covering sieve is a Nisnevich covering sieve on* CAlg^{cn}*.*
(ii) A Zariski covering sieve and a Nisnevich covering sieve are covering sieves on the flat topology on CAlg^{cn}*.*

Proof. Let R be a connective $\mathbb{E}_\infty$-ring and $f_1, \cdots, f_n$ elements in $\pi_0 R$ which generate the unit ideal of R. To prove (i), we show that each $R \to R[f_i^{-1}]$ is flat and $\{\pi_0 R \to \pi_0 R[f_i^{-1}]\}$ is an ordinary Nisnevich covering of $\pi_0 R$. Since each $R \to R[f_i^{-1}]$ is flat by [13, Remark 2.9] and $\{\pi_0 R \to \pi_0 R[f_i^{-1}]\}$ is an ordinary Zariski covering of $\pi_0 R$, so that it is a Nisnevich covering of $\pi_0 R$ [11, Remark 1.13].

To show (ii), we show that any collection of morphisms $R \to R_a$ such that their underlying maps of commutative rings $\pi_0 R \to \pi_0 R_a$ determine a Nisnevich covering, $R \to \prod_\alpha R_\alpha$ is faithfully flat. Note that we can assume that this Nisnevich covering is a finite collection of morphisms [11, Remark 1.6].

Since each $R \to R_\alpha$ is étale, it is flat. It follows that $R \to \prod_\alpha R_\alpha$ is also flat. By [11, Remark 1.12], $\pi_0 R \to \pi_0(\prod_\alpha R_\alpha) \cong \prod_\alpha \pi_0 R_\alpha$ is faithfully flat. □

3. Group completion in an ∞-topos

Definition 3.1 (cf. [9] Definition 2.4.2.1, Definition 5.2.6.2)**.** Let $\mathcal{C}$ be an ∞-category with finite products, and $\mathcal{O}^{\otimes}$ be the ∞-operad $\mathcal{A}ss^{\otimes}$ (resp. $\mathrm{N}_{\Delta}(\mathcal{F}\mathrm{in}_{*})$) given in [9, Definition 4.1.1.3] (resp. given by taking the simplicial nerve of [9, Notation 2.0.0.2]).

(i) An ∞-monoid (resp. commutative ∞-monoid) in $\mathcal{C}$ is a functor $M : \mathcal{O}^{\otimes} \to \mathcal{C}$ such that the morphism $M(\langle n \rangle) \to M(\langle 1 \rangle)$ induced by the inert maps $\sigma^{i} : \langle n \rangle \to \langle 1 \rangle$ induces an equivalence $M(\langle n \rangle) \simeq M(\langle 1 \rangle)^{n}$.

(ii) An ∞-monoid is an ∞-group if its image in $h\mathcal{C}$ is a group object.

Remark 3.2. In terminology of [6], a commutative ∞-monoid is called an $\mathbb{E}_{\infty}$-monoid, and a commutative ∞-group is called an $\mathbb{E}_{\infty}$-group by [6, Proposition 1.1].

Recall that the concepts of ∞-monoid and ∞-group are defined in Definition 3.1. In this section, we treat the several adjoint functors on ∞-categories, and we characterize the group completion functor.

3.1. **The group completion functor.**

Definition 3.3. Let $\mathcal{C}$ be an ∞-category with finite products as in Definition 3.1.

(i) In this section, according to the notation in [6, Definition 1.2], we denote by $Mon(\mathcal{C})$ an ∞-category of commutative ∞-monoids in $\mathcal{C}$ and by $Gp(\mathcal{C})$ an ∞-category of commutative ∞-groups in $\mathcal{C}$ [6, Proposition 1.1].

(ii) We say that a functor $Mon(\mathcal{C}) \to Gp(\mathcal{C})$ is a group completion on $\mathcal{C}$ if it is the left adjoint to the inclusion functor $Gp(\mathcal{C}) \to Mon(\mathcal{C})$.

(iii) Let $\mathcal{S}$ be the ∞-category of spaces. As a special case, if we take a symmetric monoidal ∞-category $\mathcal{S}$, we also use the notations $nMon(\mathcal{S})$ and $nGp(\mathcal{S})$ ∞-categories of not-necessary commutative ∞-monoids and not-neccesary commutative ∞-groups in $\mathcal{C}$ respectively. We also say that a functor $nMon(\mathcal{S}) \to nGp(\mathcal{S})$ is a group completion on $\mathcal{S}$ if it is the left adjoint to the inclusion functor $nGp(\mathcal{S}) \to nMon(\mathcal{S})$.

By the universal property of adjoint functors, the group completion functor is uniquely determined up to equivalence [8, Remark 5.2.2.2].

Definition 3.4 (cf. [9] Lemma 2.4.5.9)**.** (i) We denote by $\mathrm{Fun}^{\prod}(\mathcal{C}, \mathcal{D})$ an ∞-category of functors which preserve the finite products.

(ii) We define the product on $\mathrm{Fun}^{\prod}(\mathcal{C}, \mathcal{D})$ by taking the objectwise product in $\mathcal{S}$, i.e., the product induced from the formation of product in $\mathcal{S}$ under the sheafification.

Lemma 3.5 ([6] Lemma 1.6)**.** *Let $\mathcal{C}$ and $\mathcal{D}$ are ∞-categories with finite products. Let $\mathrm{Fun}^{\prod}(\mathcal{C}, \mathcal{D})$ be an ∞-category of functors which preserve the finite products. Then, we*

have

$$Mon(\mathrm{Fun}^{\prod}(\mathcal{C},\, \mathcal{D})) \simeq \mathrm{Fun}^{\prod}(\mathcal{C},\, Mon(\mathcal{D})),$$

and

$$Gp(\mathrm{Fun}^{\prod}(\mathcal{C},\, \mathcal{D})) \simeq \mathrm{Fun}^{\prod}(\mathcal{C},\, Gp(\mathcal{D})).$$

□

Definition 3.6 (cf. [8] 6.1.2.7)**.** Let $\mathcal{C}$ be an ∞-topos. For an ∞-monoid object G, there exists a colimit of the simplicial homotopy diagram by [9, Lemma 5.2.2.6]

$$\cdots \quad G \times G \times G \quad G \times G \quad G \quad 1.$$

where the face map is given by the multiplication on G and the degeneracy map is given by the unit morphism. We define BG is a colimit of the simplicial homotopy diagram.

Remark 3.7. Let $\mathcal{S}$ be the symmetric cartesian monoidal ∞-category of spaces. In terminology of topology, an object M in $\mathcal{S}$ is grouplike if $\pi_0 M$ is a group object in $h\mathcal{S}$ [9, Example 5.2.6.4]. A grouplike object in $\mathcal{S}$ is just an ∞-group in $\mathcal{S}$ by Definition 3.1.

If $\mathcal{C}$ is a symmetric cartesian monoidal model category, an ∞-monoid object can be regarded as an associative algebra object by [9, Proposition 2.4.2.5].

3.2. **Characterization of the group completion on $\mathcal{S}$.** Let $\mathcal{S}$ be the ∞-category of spaces, and $\mathcal{S}_*$ the pointed ∞-category of spaces (cf. [9, Notation 5.2.6.11]).

For a space X, we define the loop space of X by $\mathrm{Map}(S^1,\, X)$, where S^1 is a simplicial circle Δ^1/Δ^0.

Let $\mathcal{S}_{*\geq 1}$ denote the full ∞-subcategory of connected spaces in $\mathcal{S}_*$. There is an equivalence from the ∞-category of connected spaces to the ∞-category of not-neccesary commutative ∞-groups in $\mathcal{S}$ obtained by the adjoint functors $B : nMon(\mathcal{S}) \rightleftarrows \mathcal{S}_* : \Omega$. There is an equivalence from the ∞-category Sp^{cn} to $Gp(\mathcal{S})$ obtained by B and Ω, which is shown in [9, Remark 5.2.6.26].

Lemma 3.8. *Let $\mathcal{S}$ be an ∞-category of spaces. Let $i : nGp(\mathcal{S}) \to nMon(\mathcal{S})$ be the forgetfull functor. With the previous notations, we have the adjunction*

$$\Omega B : nMon(\mathcal{S}) \rightleftarrows nGp(\mathcal{S}) : i,$$

where i is the forgetfull functor. In other words, ΩB is the group completion on $\mathcal{S}$.

□

3.3. **Characterization of the group completion on** $\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})$**.** In this subsection, we characterize the group completion functor on $\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})$ by using Ω and B.

Definition 3.9. (i) In this paper, we will denote by ΩB the left adjoint functor given by $\Omega B : nMon(\mathcal{S}) \rightleftarrows nGp(\mathcal{S})$ in Lemma 3.8.

(ii) We also denote by ΩB the functor $\mathrm{Fun}((\mathrm{CAlg}^{\mathcal{G}})^{op}, Mon(\widehat{\mathcal{S}})) \to \mathrm{Fun}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})$ which sends a presheaf F to a presheaf given by $R \mapsto \Omega B(F(R))$.

We will give an explicit description of the group completion functor for the ∞-category of presheaves which preserves finite products.

Definition 3.10. We define a functor

$$\Omega B^{\prod} : Mon(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) \to Gp(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}}))$$

by the restriction of ΩB, i.e., $\Omega B^{\prod}$ is the functor which satisfies the following commutative diagram

$$\begin{array}{ccc} Mon(\mathrm{Fun}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) & \xrightarrow{\Omega B} & Gp(\mathrm{Fun}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) \\ \uparrow & & \uparrow \\ Mon(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) & \xrightarrow{\Omega B^{\prod}} & Gp(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) \end{array},$$

where the vertical morphisms are inclusions.

Lemma 3.11. *The functor* $\Omega B^{\prod}$ *gives the group completion on* $\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})$ *up to equivalences.*

Proof. Let $\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})$ be an ∞-category of presheaves on $\mathrm{CAlg}^{\mathcal{G}}$ which preserve the finite products. This ∞-category has the pointwise finite products. By Lemma 3.5, we have the equivalences $Mon(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) \simeq \mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, Mon(\widehat{\mathcal{S}}))$ and $Gp(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) \simeq \mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, Gp(\widehat{\mathcal{S}}))$.

Therefore, the group completion functor on $\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})$ is determined by the values in the ∞-category $\mathcal{S}$ of spaces. The problem is reduced to the group completion functor $nMon(\mathcal{S}) \to nGp(\mathcal{S})$ for the ∞-category $\mathcal{S}$ of spaces, which is ΩB by Lemma 3.8. □

3.4. **Characterization of the group completion on** $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$**.**

Definition 3.12 (cf. [13] Proposition 1.15)**.** We define the product on $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$ by the pointwise product, i.e., the product induced from the formation of product in $\mathcal{S}$ under the sheafification.

Since a sheaf preserves finite limits, the pointwise product on $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$ is the restriction of the product on $\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})$ in Definition 3.4.

Lemma 3.13. *We have* $\mathcal{S}hv_{Gp(\widehat{\mathcal{S}})}(\mathrm{CAlg}^{\mathcal{G}}) \subset Gp(\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}}))$.

Proof. By definition, the objectwise products of presheaves becomes pointwise products of sheaves after sheafification. Since the sheafification is left exact [8, Definition 5.3.2.1], it commutes with the finite products. Therefore, we have $\mathcal{S}hv_{Gp(\widehat{\mathcal{S}})}(\mathrm{CAlg}^{\mathcal{G}}) \subset Gp(\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}}))$. □

Definition 3.14. We define the functor

$$\Omega B^{\mathcal{G}} : Mon(\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})) \to Gp(\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$$

by the composition of the inclusion $i' : Mon(\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})) \to Mon(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}}))$ with

$$Mon(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) \to Gp(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) \to Shv_{Gp(\widehat{\mathcal{S}})}(\mathrm{CAlg}^{\mathcal{G}}),$$

where the first functor is the functor which is induced from the pointwise group completion $\Omega B^{\prod} : Mon(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}})) \to Gp(\mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, \widehat{\mathcal{S}}))$, and the second functor is obtained by the equivalence in Lemma 3.5 and the sheafification functor $(\tilde{-}) : \mathrm{Fun}^{\prod}((\mathrm{CAlg}^{\mathcal{G}})^{op}, Gp(\widehat{\mathcal{S}})) \to Shv_{Gp(\widehat{\mathcal{S}})}(\mathrm{CAlg}^{\mathcal{G}})$. Note that, we have $\mathcal{S}hv_{Gp(\widehat{\mathcal{S}})}(\mathrm{CAlg}^{\mathcal{G}}) \subset Gp(\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}}))$ by Lemma 3.13.

Proposition 3.15. *Let* $\mathrm{CAlg}^{\mathcal{G}}$ *be the* ∞*-category equipped with the Grothendieck topology which is defined in Section 2. Then,* $\Omega B^{\mathcal{G}}$ *is the group completion on* $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$.

Proof. By the adjunction in Lemma 3.8 and the definition of $\Omega B^{\mathcal{G}}$, we have

$$\mathrm{Map}(\Omega B^{\mathcal{G}} F,\, G) \simeq \mathrm{Map}((-)^{\simeq} \circ \Omega B \circ i'(F),\, G) \simeq \mathrm{Map}(\Omega B i F,\, iG) \simeq \mathrm{Map}(F,\, iG).$$

□

4. Classifying sheaf of GL and projective modules of finite rank

4.1. **The affine spectral scheme GL_n.** For a simplicial set S and its vertexes x and y, recall that a simplicial set $\mathrm{Map}_S(x,\, y)$ is defined as follows. This construction is due to Joyal.

Definition 4.1. [cf. [8] 1.2.2.2, Corollary 4.2.1.8] Let S be a simplicial set and S^{Δ^1} a simplicial set which sends $[n]$ to $\mathrm{Hom}_{\mathrm{Set}_\Delta}(\Delta^1 \times \Delta^n,\, S)$. Let $s, t : [0] \to [1]$ be maps defined by $s(0) = 0$ and $t(0) = 1$.

(i) Take vertexes $x, y \in S$. The mapping space $\mathrm{Map}_S(x,\, y)$ from x to y is defined by the pullback

$$\begin{array}{ccc} \mathrm{Map}_S(x,\, y) & \longrightarrow & S^{\Delta^1} \\ \downarrow & & \downarrow{\scriptstyle (s,t)} \\ * & \xrightarrow{(x,y)} & S \times S, \end{array}$$

where the morphism (s,t) is induced by s and t, e.g., it sends $\mathrm{Hom}_{\mathrm{Set}_\Delta}(\{0\}\times\Delta^n,\ S)$ to the n-simplices S_n of the first factor and $\mathrm{Hom}_{\mathrm{Set}_\Delta}(\{1\}\times\Delta^n,\ S)$ to the n-simplices S_n of the second factor, and the morphism (x,y) sends $*$ to $(x,y)\in S\times S$.

(ii) If S is an ∞-category, $\mathrm{Map}_S(x,\ y)$ becomes a Kan complex by [8, Proposition 1.2.2.3]. For an ∞-category S and objects $x,y\in S$, we say that $\mathrm{Map}_S(x,\ y)$ is the mapping space between x and y.

Definition 4.2 ([9] Notation 3.1.3.8)**.** Let R be an $\mathbb{E}_\infty$-ring, and CAlg_R the ∞-category of R-algebras.

(i) We define a functor $\mathrm{Sym}_R : \mathrm{Mod}_R \to \mathrm{CAlg}_R$ the left adjoint of the forgetful functor $\mathrm{CAlg}_R \to \mathrm{Mod}_R$ which sends an R-algebra S to an R-module S.

(ii) For a free R-module $R^{\oplus n^2}$ of rank n^2, $\pi_0\mathrm{Sym}_R R^{\oplus n^2}$ is isomorphic to the polynomial ring $(\pi_0 R)[x_{11},\cdots,x_{nn}]$ over $\pi_0 R$. Let us denote $\mathrm{Sym}_R R^{\oplus n^2}$ by $R\{x_{11},\cdots,x_{nn}\}$, where each x_{ij} is the representative of the indeterminate of the polynomial ring over $\pi_0 R$.

Definition 4.3 ([9] Section 4.7.2)**.** For an object X in a symmetric monoidal ∞-category, an endomorphism object $\mathrm{End}(X)$ is an object equipped with the evaluation morphism $e : \mathrm{End}(X)\otimes X \to X$ which induces a weak homotopy equivalence $\mathrm{Map}(Y\otimes X,\ X)\simeq \mathrm{Map}(Y,\ \mathrm{End}(X))$ for every Y.

Let R be an $\mathbb{E}_\infty$-ring. We have the endomorphism object of $R^{\oplus n}$ in Mod_R. Let us denote it by $\mathrm{End}_R(R^n)$.

The following lemma is explained in [9, Remark 7.1.2.2].

Lemma 4.4 ([9] Remark 7.1.2.2)**.** *We have an isomorphism*

$$\pi_*\mathrm{End}_R(R^{\oplus n}) \cong \pi_*\mathrm{Map}_R(R^{\oplus n},\ R^{\oplus n})$$

□

The object $R^{\oplus n^2}$ of Mod_R satisfies the universal property of endomorphism, we have an equivalence $R^{\oplus n^2}\simeq \mathrm{End}_R(R^{\oplus n})$. Since we have $R^{\oplus n^2}\simeq \mathrm{End}_R(R^{\oplus n})$, we have an equivalence $\mathrm{Sym}_R\mathrm{End}_R(R^{\oplus n})\simeq R\{x_{11},\cdots,x_{nn}\}$ by Definition 4.2 (ii).

Definition 4.5. Let R be an $\mathbb{E}_\infty$-ring. Let us denote $\mathrm{Spec}^{Zar}\mathrm{Sym}_R\mathrm{End}_R(R^{\oplus n})$ by $M_{n,R}$ in $\mathcal{S}hv(\mathrm{CAlg}^{Zar})$.

(i) Under the equivalence $\mathrm{Sym}_R\mathrm{End}_R(R^{\oplus n})\simeq R\{x_{11},\cdots,x_{nn}\}$, we define the element $(det)\in\pi_0 M_{n,R}$ by the determinant relation $\Sigma_{\tau\in S_n} sgn(\tau)\prod x_{i,\tau(i)}$ of x_{ij}s.

(ii) We define an affine scheme $GL_{n,R}$ by inverting the determinant element of $M_{n,R}$.

(iii) In the case that the base scheme R is the sphere spectrum $\mathbb{S}$, we denote $M_{n,R}$ and $GL_{n,\mathbb{S}}$ by M_n and GL_n.

Let S be an R-algebra. We denote by $M_{n,R}(S)$ and $GL_{n,R}(S)$ the mapping spaces $\mathrm{Map}_{\mathrm{CAlg}_R}(\mathrm{Sym}_R\mathrm{End}_R(R^{\oplus n}), S)$ and $\mathrm{Map}_{\mathrm{CAlg}_R}(\mathrm{Sym}_R\mathrm{End}_R(R^{\oplus n})[(det)^{-1}], S)$ respectively.

Remark 4.6. Since GL_n is corepresented by an $\mathbb{E}_\infty$-ring, it is flat sheaf by [13, VII, Theorem 5.15], so that it is already a Nisnevich sheaf. We also use the notation GL_n for the image in $\mathcal{S}hv(\mathrm{CAlg}^{Nis})$ under the sheafification.

4.2. **The equivalence $GL_n(R) \simeq Aut(R^n)$ as ∞-groups.**

Definition 4.7. We define the space $Aut_R(S^n)$ by the following pullback of simplicial sets

$$\begin{array}{ccc} Aut_R(S^n) & \longrightarrow & \mathrm{Map}_{\mathrm{Mod}_R}(S^n, S^n) \\ \downarrow & & \downarrow{\scriptstyle \pi_0} \\ (\pi_0\mathrm{Map}_{\mathrm{Mod}_R}(S^n, S^n))^\times & \longrightarrow & \pi_0\mathrm{Map}_{\mathrm{Mod}_R}(S^n, S^n), \end{array}$$

where we regard $(\pi_0\mathrm{Map}_{\mathrm{Mod}_R}(R^n, R^n))^\times$ and $\pi_0\mathrm{Map}_{\mathrm{Mod}_R}(R^n, R^n)$ as constant simplicial sets and $(\pi_0\mathrm{Map}_{\mathrm{Mod}_R}(R^n, R^n))^\times$ is the invertible objects in $\pi_0\mathrm{Map}_{\mathrm{Mod}_R}(R^n, R^n)$.

Remark 4.8. Note that the constant simplicial sets which appear in the diagram of Definition 4.7 are also Kan complexes since the homotopy set of the mapping space is made from the same mapping space by replacing 1-simplices with isomorphism. (Note that any 1-simplex in a Kan complex is invertible.)

Proposition 4.9. *For an R-algebra S, we have an equivalence $GL_{n,R}(S) \simeq Aut_R(S^n)$ as ∞-groups in $\mathcal{S}$, which is functorial with respect to R-algebra S.*

Proof. Note that the right vertical morphism π_0 in the diagram in Definition 4.7 is a Kan fibration.

Since $GL_{n,R}(S)$ is formulated by the following pullback of simplicial sets

$$\begin{array}{ccc} GL_{n,R}(S) & \longrightarrow & M_{n,R}(S) \\ \downarrow & & \downarrow{\scriptstyle \pi_0} \\ (\pi_0 M_{n,R}(S))^\times & \longrightarrow & \pi_0 M_{n,R}(S), \end{array} \tag{4.1}$$

where we regard $\pi_0 M_{n,R}(S)$ as a constant simplicial set and $(\pi_0 M_{n,R}(S))^\times$ is the invertible objects in $\pi_0 M_{n,R}(S)$, and by the coglueing lemma (cf. [8] A.2.4.3), it suffices to construct the two morphisms $M_{n,R}(S) \to \mathrm{Map}_{\mathrm{Mod}_R}(S^n, S^n)$ and $\pi_0 M_{n,R}(S) \to \pi_0\mathrm{Map}_{\mathrm{Mod}_R}(S^n, S^n)$ which preserve the multiplication and show that the following diagram is commutative:

$$\begin{array}{ccccc} (\pi_0 M_{n,R}(S))^\times & \longrightarrow & \pi_0 M_{n,R}(S) & \underset{\pi_0}{\longleftarrow} & M_{n,R}(S) \\ \downarrow & & \downarrow & & \downarrow \\ (\pi_0\mathrm{Map}_{\mathrm{Mod}_R}(S^n, S^n))^\times & \longrightarrow & \pi_0\mathrm{Map}_{\mathrm{Mod}_R}(S^n, S^n) & \underset{\pi_0}{\longleftarrow} & \mathrm{Map}_{\mathrm{Mod}_R}(S^n, S^n). \end{array} \tag{4.2}$$

Note that objects in the diagram (4.1) are Kan complexes by the same reason of Remark 4.8

An element of $M_{n,R}(S)$ is a morphism $\mathrm{Sym}_R\mathrm{End}_R(R^{\oplus n}) \to S$ of R-algebras for an R-algebra S. By the adjointness of Sym_R, the morphism of R-algebras is corresponding to a morphism of R-modules

$$R^{\oplus n^2} \simeq \mathrm{End}_R(R^{\oplus n}) \to S.$$

On the other hand, we have an equivalence $R^{\oplus n^2} \simeq \mathrm{End}_R(R^{\oplus n})$ obtained by evaluating the each factor of R^n. Therefore, if we regard $S \otimes_R R^{\oplus n^2}$ as an S-module, the above morphism corresponds to a morphism $S \otimes_R R^{\oplus n^2} \to S$ of S-modules. This gives an identification $M_{n,R}(S) \simeq \mathrm{End}_S(S^{\oplus n})$ as S-modules.

To show that $GL_{n,R}(S) \to Aut_R(S^n)$ is a morphism of ∞-groups, we fix the choice $\mathrm{Map}_{\mathrm{Mod}_S}(S^{\oplus n^2}, S) \simeq \mathrm{Map}_{\mathrm{Mod}_S}(S, S)^{\oplus n^2} \simeq \mathrm{Map}_{\mathrm{Mod}_S}(S^n, S^n)$ of the second equivalence. Then, by the composition of S-module endomorphisms on $S^{\oplus n}$, $\mathrm{End}_S(S^{\oplus n})$ has a canonical ∞-monoid structure for each S defined in Definition 3.1, the spectral scheme $M_{n,R}$ is an ∞-monoid.

We can identify the discrete group $\pi_0 M_{n,R}(S)^\times$ with the class of $\pi_0 S$-algebra morphisms $\pi_0\mathrm{Sym}_R\mathrm{End}_R(R^{\oplus n})[det^{-1}] \to \pi_0 S$. Here, det is the determinant element given by the determinant relation of $x_{ij}s$ in $\pi_0 R[x_{11}, \cdots, x_{nn}]$. By ordinary theory of affine group schemes, these morphisms corresponds to $(\pi_0\mathrm{End}_S(S^{\oplus n}))^\times$ by the same choice of isomorphism $\mathrm{Hom}(\pi_0 S, \pi_0 S)^{\oplus n^2} \simeq \mathrm{Hom}(\pi_0 S^n, \pi_0 S^n)$. Since 2-morphisms are invertible, this induces an equivalence between $Aut_R(S^n)$ and $\mathrm{Sym}_R\mathrm{End}_R(R^{\oplus n})[det^{-1}] \to S$ by the above pullback.

Since we fix the choice of equivalences in the proof, the affine scheme $GL_{n,R}$ is an ∞-group scheme with respect to the monoid structure of $\mathrm{Map}_{\mathrm{Mod}_R}(R^n, R^n)$. □

4.3. **The $\widehat{\mathrm{Cat}_\infty}$-valued functor** $(nProj)$**.** In this subsection, the functor $(nProj) : (\mathrm{CAlg}^{cn})^{op} \to \widehat{\mathrm{Cat}_\infty}$ defined by sending an $\mathbb{E}_\infty$-ring R to the maximal Kan complex of ∞-category $\mathrm{Mod}_R^{\infty proj}$ of projective modules of finite rank is a sheaf.

Definition 4.10 ([10] Definition 2.6.14)**.** Let P be a property for objects (A, M) in an ∞-category $\mathrm{CAlg}^{cn} \times_{\mathrm{CAlg}} \mathrm{Mod}$. We say that P is local for the flat topology if the following conditions are satisfied:

(i) Let $f : A \to B$ be a flat morphism of connective $\mathbb{E}_\infty$-rings, and M an A-module. If (A, M) has the property P, $(B, B \otimes_A M)$ has the property P. If f is faithfully flat, the converse holds.

(ii) For any finite collection (A_i, M_i) of the objects in $\mathrm{CAlg}^{cn} \times_{\mathrm{CAlg}} \mathrm{Mod}$ such that each (A_i, M_i) has the property P, the product $(\prod A_i, \prod M_i)$ has the property P.

Theorem 4.11 ([13] Corollary 6.13, Lemma 6.17)**.** *A functor* $(\mathrm{CAlg}^{cn})^{op} \to \widehat{\mathrm{Cat}_\infty}$ *given by* $R \mapsto \mathrm{Mod}_R$ *is a sheaf.*

Lemma 4.12. *Let* $(nProj) : (\mathrm{CAlg}^{cn})^{op} \to \widehat{\mathrm{Cat}_\infty}$ *be a functor given by* $R \mapsto \mathrm{Mod}_R^{nproj}$*. Then, the functor* $(nProj)$ *is a sheaf with respect to flat topology.*

Proof. Since a functor $(\mathrm{CAlg}^{cn})^{op} \to \widehat{\mathrm{Cat}_\infty}$ given by $R \mapsto \mathrm{Mod}_R$ is already a sheaf By Theorem 4.11, it suffices to check that the projective modules of rank n satisfy the condition of Definition 4.10 (i) and (ii).

By [10, Proposition 2.6.15 (1), (6), (9)], the condition of finitely generated projective is flat local property. Since the tensor product preserves rank, the projective modules of rank n satisfy the condition of Definition 4.10 (i).

We will check the condition (ii) of Definition 4.10. Assume that $(A_i,\, M_i)$ is a pair such that A_i is connective $\mathbb{E}_\infty$-ring and M_i is a projective A_i-module of rank n for $1 \leq i \leq m$. To show that $\prod_i M_i$ is a projective $\prod_i A_i$-module of rank n, it suffices to show that there exists a finite set $\{g_a\}_a$ of objects in $\prod_i A_i$ such that each $\prod_i M_i[g_a^{-1}]$ is a free $\prod_i A_i[g_a^{-1}]$-module of rank n. We choose such $\{g_a\}_a$ as follows.

For each i, we take $f_{i1}, \cdots, f_{ik} \in \pi_0 A_i$ such that $(M_i)[f_{il}^{-1}]$ is a free $A_i[f_{il}^{-1}]$-module of rank n for $1 \leq l \leq k$. Since $\prod_i A_i \to A_i$ is flat and the essential image is generated by the form $\prod(A_i \otimes_A A_j) \otimes_{A_j} M$ and $A_i \otimes_A A_j \simeq 0$, $A_j \otimes_{\prod_i A_i} \prod_i M_i \simeq M_j$.

From this, we have the equivalence $\prod_i A_i[f_{il}^{-1}] \otimes_{\prod_i A_i} \prod_i M_i \simeq \prod_i M_i[f_{il}^{-1}]$, where $\prod_i M_i[f_{il}^{-1}]$ is regarded as a $\prod_i A_i[f_{il}^{-1}]$-module and is free of rank n. □

Corollary 4.13. *(nProj) is a Zariski and Nisnevich sheaf.*

Proof. It follows from Lemma 2.7 and Lemma 4.12. □

Definition 4.14 ([8] Theorem 3.1.5.1)**.** Let $\widehat{\mathcal{S}}$ be the ∞-category of spaces and $\widehat{\mathrm{Cat}_\infty}$ the ∞-category of ∞-categories defined in Definition 2.3 after enlarging the universe respectively. We regard the Kan complexes as ∞-categories, so that we have an inclusion functor $i : \widehat{\mathcal{S}} \to \widehat{\mathrm{Cat}_\infty}$.

Since the inclusion preserves small colimits, by [8, Corollary 5.5.2.9], there is an adjunction

$$i : \widehat{\mathcal{S}} \rightleftarrows \widehat{\mathrm{Cat}_\infty}(-)^{\simeq}. \tag{4.3}$$

Note that the right adjoint $(-)^{\simeq}$ is given by taking the maximal ∞-groupoid.

Lemma 4.15. *The right adjoint in (4.3) induces a functor from* $\widehat{\mathrm{Cat}_\infty}$*-valued sheaves to* $\widehat{\mathcal{S}}$*-valued sheaves.*

Proof. Since $(-)^{\simeq}$ is a right adjoint, it preserves limits. Especially, it preserves the limits of simplicial diagrams in Proposition 2.6. Therefore, $(-)^{\simeq}$ sends $\widehat{\mathrm{Cat}_\infty}$-valued sheaves to $\widehat{\mathcal{S}}$-valued sheaves. □

4.4. **Comparison between $B(GL_n(R))$ to $(\mathrm{Mod}_R^{nfree})^{\simeq}$.** Recall that the notion of classifying object from Definition 3.6. We denote by $B^{\mathcal{G}}$ the classifying functor on $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$. Acording to Remark 4.6, we have GL_n in $\mathcal{S}hv(\mathrm{CAlg}^{\mathcal{G}})$.

Definition 4.16. (i) We use the notation $B(GL_n(R))$ for the classifying space of the value $GL_n(R)$ in $\mathcal{S}$, and $(B^{\mathcal{G}}GL_n)(R)$ for the value at R of the classifying sheaf of GL_n in $\mathcal{S}hv(\mathrm{CAlg}^{\mathcal{G}})$.

(ii) Let $(\mathrm{Mod}_{(-)}^{nfree})^{\simeq} : (\mathrm{CAlg}^{cn})^{op} \to \widehat{\mathcal{S}}$ be a functor which sends R to the ∞-groupoid $(\mathrm{Mod}_R^{nfree})^{\simeq}$ of free R-modules of rank n.

(iii) Let $B(GL_n(-)) : (\mathrm{CAlg}^{cn})^{op} \to \widehat{\mathcal{S}}$ be a functor given by $R \mapsto B(GL_n(R))$.

Note that $GL_n(R) \simeq Aut_R(R^{\oplus n})$, and $B(GL_n(R))$ is not equal to $(B^{\mathcal{G}}GL_n)(R)$.

Remark 4.17. Apparently, the classifying sheaf $B^{\mathcal{G}}GL_n$ of GL_n in $\mathcal{S}hv(\mathrm{CAlg}^{\mathcal{G}})$ in Definition 4.16 depends the Grothendieck topology on $\mathrm{CAlg}^{\mathcal{G}}$, but consequently, we show that it is equivalent to the flat sheaf $(nProj)$ in Proposition 4.21 below, $B^{\mathcal{G}}GL_n$ in Zariski topology is equivalent to that in Nisnevich topology.

Now, we characterize a functor in Definition 4.16(ii) with a functor in Definition 4.16(iii).

Lemma 4.18. *Let us regard $(\mathrm{Mod}_R^{nfree})^{\simeq}$ as a simplicial set, and take a vertex $x \in (\mathrm{Mod}_R^{nfree})^{\simeq}$.*

(i) We have a morphism from $Aut_R(x)$ to $(\mathrm{Mod}_R^{nfree})^{\simeq}$ which is compatible with the face maps and the degeneracy maps, i.e., a morphism of simplicial sets.

(ii) Under the morphism in (i), we have a morphism $B(Aut_R(x)) \to (\mathrm{Mod}_R^{nfree})^{\simeq}$ of simplicial sets.

Proof. If we have $x = y$ in the setting in Definition 4.1, we have a morphism $\mathrm{Map}(x,\, x) \to X$ of simplicial set by restricting the projection (s, t) on the first or the second factors. Thus, we have a morphism from $\mathrm{Map}(x, x)$ to X which is compatible with the face maps and the degeneracy maps. By composing the projection from n-times product $\mathrm{Map}(x, x)^n$ to $\mathrm{Map}(x, x)$ with $\mathrm{Map}(x, x) \to X$, we have $\mathrm{Map}(x, x)^n \to X$ which is also compatible with the face maps and the degeneracy maps.

For (i), by applying the above argument to the simplicial set Mod_R^{nfree} and a vertex $x \in \mathrm{Mod}_R^{nfree}$, we obtain the morphisms $\mathrm{Map}_{\mathrm{Mod}_R}(x,\, x) \to \mathrm{Mod}_R^{nfree}$ and $\mathrm{Map}_{\mathrm{Mod}_R}(x,\, x)^n \to \mathrm{Mod}_R^{nfree}$ which is compatible with the face and degeneracy maps, i.e., morphisms of simplicial sets.

Recall that $Aut_R(x)$ defined in Definition 4.7 is a simplicial subset of the mappling space $\mathrm{Map}_{\mathrm{Mod}_R}(x,x)$. Then, the induced morphisms $Aut_R(x) \to \mathrm{Map}_{\mathrm{Mod}_R}(x,\, x) \to \mathrm{Mod}_R^{nfree}$ factors through the maximal Kan complex $Aut_R(x) \to (\mathrm{Mod}_R^{nfree})^{\simeq}$. From this, we have the morphism $Aut_R(x)^n \to (\mathrm{Mod}_R^{nfree})^{\simeq}$ which is compatible with the face and degeneracy maps.

For (ii), by the above argument, we have the morphism $Aut_R(x)^n \to (\mathrm{Mod}_R^{nfree})^{\simeq}$ which is compatible with the face and degeneracy maps, i.e., morphism of simplicial set, from the each factor of $B(Aut_R(x))$ to $(\mathrm{Mod}_R^{nfree})^{\simeq}$. Therefore, it induces a morphism $B(Aut_R(x)) \to (\mathrm{Mod}_R^{nfree})^{\simeq}$ of simplicial sets. □

Proposition 4.19. *A functor in Definition 4.16(ii) is characterized by a functor in Definition 4.16(iii).*

Let $(\mathrm{Mod}_{(-)}^{nfree})^{\simeq} : (\mathrm{CAlg}^{cn})^{op} \to \widehat{\mathbb{S}}$ *be a functor which sends* R *to the* ∞*-groupoid* $(\mathrm{Mod}_R^{nfree})^{\simeq}$ *of free* R*-modules of rank* n*. Let* $B(GL_n(-)) : (\mathrm{CAlg}^{cn})^{op} \to \widehat{\mathbb{S}}$ *be a functor given by* $R \mapsto B(GL_n(R))$*. Then, as a functor,* $B(GL_n(-)) \simeq (\mathrm{Mod}_{(-)}^{nfree})^{\simeq}$.

Proof. By taking $x = y = R^n$ in Lemma 4.18, we have a morphism from $B(GL_n(R)) \to (\mathrm{Mod}_R^{nfree})^{\simeq}$ for each $R \in (\mathrm{CAlg}^{cn})^{op}$.

By composing a morphism $B(GL_n(R)) \simeq BAut_{\mathbb{S}}(R^n)$ obtained by Proposition 4.9 with $B(Aut_{\mathbb{S}}(R)) \to (\mathrm{Mod}_R^{nfree})^{\simeq}$ obtained by Lemma 4.18, we have a morphism $B(GL_n(R)) \to (\mathrm{Mod}_R^{nfree})^{\simeq}$. It suffices to show that a morphism $B(GL_n(R)) \to (\mathrm{Mod}_R^{nfree})^{\simeq}$ is an equivalence and functorial with respect to R.

By the construction, we identify $B(GL_n(R))$ with a full ∞-subgroupoid of $(\mathrm{Mod}_R^{nfree})^{\simeq}$ such that the object is only $R^{\oplus n}$ and the class of morphisms is identified with $GL_n(R) \simeq Aut_{\mathbb{S}}(R)$ by Proposition 4.9, which is fully faithfull functorial assignment with respect to R.

We show that it is essentially surjective. It suffices to show that $(\mathrm{Mod}_R^{nfree})^{\simeq}$ is connected as a simplicial set. This is obvious since, for $M \in \mathrm{Mod}_R^{nfree}$, we have $M \simeq R^{\oplus n}$. □

4.5. **A natural transformation from $B(GL_n(-))$ to $(B^{\mathcal{G}}GL_n)(-)$.** We construct a natural transformation from $B(GL_n(-))$ to $(B^{\mathcal{G}}GL_n)(-)$ by using the comparison between the sheaf of projective modules of rank n and the sheafification of the functor of free modules of rank n.

For each R, we have the following colimit in $\widehat{\mathbb{S}}$:

$$GL_n(R) \rightrightarrows * \longrightarrow B(GL_n(R)).$$

Note that the functor corresponds to this cofiber is not a sheaf since the second condition of [10, Definition 2.6.14 (1)] fails. According to the notation in Definition 2.4, we write $B(\widetilde{GL_n}(-))$ for the sheafification of $B(GL_n(-))$.

Lemma 4.20. *We have the following cofiber sequence*

$$GL_n \rightrightarrows * \longrightarrow B(\widetilde{GL_n(-)})$$

in $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$.

Proof. Since GL_n is corepresented by an $\mathbb{E}_\infty$-ring, it is flat sheaf by [13, VII, Theorem 5.15], so that it is a Zariski (resp. Nisnevich) sheaf. Since sheafification is left adjoint, it commutes with the cofiber sequence. □

For each $A \to R$ in CAlg^{cn}, the natural inclusions induce the commutative diagram

$$\begin{array}{ccc} \mathrm{Mod}_A^{nfree} & \longrightarrow & \mathrm{Mod}_A^{nproj} \\ \downarrow & & \downarrow \\ \mathrm{Mod}_R^{nfree} & \longrightarrow & \mathrm{Mod}_R^{nproj}, \end{array}$$

in $\widehat{\mathrm{Cat}_\infty}$, so that we have a natural transformation

$$\mathrm{Mod}_{(-)}^{nfree} \to \mathrm{Mod}_{(-)}^{nproj}. \tag{4.4}$$

By the adjointness of sheafification, we have a morphism $B(\widetilde{GL_n(-)}) \to (nProj)^{\simeq}$ of $\widehat{\mathcal{S}}$-valued sheaves.

4.6. **An equivalence $B^{\mathcal{G}}GL_n \simeq (nProj)^{\simeq}$.**

Proposition 4.21. *The morphism (4.4) in the previous subsection gives an equivalence* $B^{\mathcal{G}}GL_n \simeq (nProj)^{\simeq}$ *in* $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$.

Proof. From adjointness of a morphism $f_{(-)} : B(\widetilde{GL_n(-)}) \to (nProj)^{\simeq}$ of $\widehat{\mathcal{S}}$-valued sheaves, we have the following homotopy commutative diagram of $\widehat{\mathcal{S}}$-valued presheaves:

$$\begin{array}{ccc} & & B(\widetilde{GL_n(-)}) \\ & \overset{L}{\nearrow} & \\ B(GL_n(-)) & \xrightarrow{\quad i \quad} & (nProj)^{\simeq}, \end{array} \qquad \text{with } f : B(\widetilde{GL_n(-)}) \to (nProj)^{\simeq},$$

where L is the morphism associated to the sheafification and i is induced from the inclusion.

Let us denote the limits of the following simplicial diagrams by $B(GL_n(A))_R$:

$$B(GL_n(A)) \rightrightarrows B(GL_n(A \otimes_R A)) \;\substack{\rightarrow\\ \rightarrow\\ \rightarrow}\; \cdots,$$

where $R \to A$ is a faithfully flat morphism. By Remark 2.5, the sheafification $B(\widetilde{GL_n(-)})$ is described by the term of $B(GL_n(-))$, i.e., it is the colimit of those $B(GL_n(A))_R$ taken over the every covering sieve $R \to A$.

For $f_R : B(\widetilde{GL_n(R)}) \to (nProj)^{\simeq}(R)$ and an object Q in $(nProj)^{\simeq}(R)$, let $(\widetilde{Q})^{\simeq}$ denote the full ∞-subgroupoid spanned by Q. We construct $h_{R,Q} : (\widetilde{Q})^{\simeq} \to B(\widetilde{GL_n(R)})$ as follows.

We take a Zariski local A of R such that an object $Q \in (nProj)^{\simeq}(R)$ is trivialized on A. Then, we naturally identified the ∞-groupoid spanned by $Q \otimes A$ with $B(GL_n(A))$, so that we have an object in $B(\widetilde{GL_n(R)})$. This is the assignment under the morphisms of ∞-groupoids. By $f \circ L \simeq i$ in the above diagram, we have $f_R \circ h_{R,Q}$ is homotopic to identity.

Conversely, for an object P in $B(\widetilde{GL_n(R)})$, we take its value on the covering sieve $R \to R$ under the morphisms associated to the colimit, and denote by P'. Then, by an extension of the coefficients of P' gives an object in $B(\widetilde{GL_n(R)})$, which is equivalent to P by the construction. This shows that $h_{R,f(P)} \circ f_R$ is homotopic to identity.

Since f_R is obviously a Kan fibration and, by the above arguments, any fiber of f_R is contractible. By [8, Lemma 4.1.3.2, Corollary 4.1.2.6], it is a weak homotopy equivalence. □

4.7. **Zariski connected $\mathbb{E}_\infty$-rings.** Next, we consider the condition on a connective $\mathbb{E}_\infty$-ring R such that any finitely generated projective R-module has finite constant rank.

Definition 4.22. (i) We say that an $\mathbb{E}_\infty$-ring is Zariski non-connected if there exists those objects $f, g \in \pi_0 R$ such that $R[f^{-1}] \otimes_R R[g^{-1}] \simeq 0$ and $R \simeq R[f^{-1}] \times R[g^{-1}]$.
(ii) We say that an $\mathbb{E}_\infty$-ring is Zariski connected if it is not Zariski non-connected.

Lemma 4.23. *The following conditions are equivalent:*

(i) R be an $\mathbb{E}_\infty$-ring such that $\pi_0 R$ has no non-trivial idempotent element.
(ii) R is Zariski connected.
(iii) Any $P \in \mathrm{Mod}_R^{proj}$ has finite constant rank.

Proof. For proving (iii) from (i), let R be an $\mathbb{E}_\infty$-ring such that $\pi_0 R$ has no non-trivial idempotent, and $P \in \mathrm{Mod}_R^{proj}$. Note that $\pi_0 P$ is a finitely generated projective $\pi_0 R$-module [9, Remark 7.2.2.20]. Then, there exists $f_1, \cdots, f_m$ which generate the unit ideal of $\pi_0 R$ such that each $\pi_0 P[f_i^{-1}]$ is free of finite rank n_i over $\pi_0 R[f_i^{-1}]$. For an ordinary commutative ring $\pi_0 R$, in this the condition (i) is equivalent to that n_is are constant, and let us denote it by n. Since finitely generated free $\pi_0 R[f_i^{-1}]$-modules $\pi_0 P[f_i^{-1}]$ can be lifted by finitely generated $R[f_i^{-1}]$-modules $P[f_i^{-1}]$ and a morphism between flat R-modules is an equivalence if and only if it induces an isomorphism on π_0 [9, Lemma 7.2.2.17], locally P has a constant rank.

Conversely, if $\pi_0 R$ has a non-trivial idempotent element, we have that the rank n_i are not constant. In this case, the above argument shows that (iii) implies (i).

We show that (i) implies (ii). Since π_0 preserves finite products, if R is Zariski non-connected, by passing to π_0 and applying the ordinary commutative ring theory, $\pi_0 R$ has a non-trivial idempotent. Conversely, if $\pi_0 R$ has a non-trivial idempotent e, it also a non-trivial idempotent of $\pi_* R$. Since the localization of an $\mathbb{E}_\infty$-ring with one element commutes with π_*, we have $\pi_*(R[e^{-1}]) \cong (\pi_* R)[e^{-1}]$ and $\pi_*(R[(1-e)^{-1}]) \cong (\pi_* R)[(1-e)^{-1}]$. Since π_* commutes with finite products, by taking π_* of $R[e^{-1}] \times R[(1-e)^{-1}]$, we conclude that R is equivalent to $R[e^{-1}] \times R[(1-e)^{-1}]$, so that R is Zariski non-connected. Thus, (ii) implies (i). □

4.8. **Comparison between $B^{\mathcal{G}}GL$ and $\mathrm{Mod}_R^{\infty proj}$.** Now, we will prove the main proposition in this section by applying the following proposition.

Proposition 4.24 ([13] Lemma 3.21)**.** *Let $\mathcal{C}$ be an ∞-topos, I an index set and $\{X_i\}_{i\in I}$ a collection of objects in $\mathcal{C}$. For every subset $J \subset I$, let $X_J \simeq \coprod_{i\in J} X_i$. Let $C \in \mathcal{C}$ be an object such that every covering of C has a finite subcovering* [13, Definition 3.1]*. Then, the canonical morphism*

$$\operatorname*{colim}_{I\subset I} \mathrm{Map}_{\mathcal{C}}(C,\, X_J) \to \mathrm{Map}_{\mathcal{C}}(C,\, X_I)$$

induces a homotopy equivalence. Here, the colimit in the left hand side is run through the all finite subsets $J \subset I$.

□

Lemma 4.25. *Let $I \subset \mathbb{N}$ be a finite index set. If R is Zariski connected, the canonical morphism*

$$\coprod_{i\in I} \mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\, B^{\mathcal{G}}GL_i) \to \mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\, \coprod_{i\in I} B^{\mathcal{G}}GL_i),$$

is an equivalence.

Proof. To check that the canonical morphism is an equivalence, it suffices to show that, for any morphism ϕ in $\mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R, \coprod_{i\in I} B^{\mathcal{G}}GL_i)$, there exists an index $j \in \mathbb{N}$ which is uniquely determined by ϕ, the morphism ϕ has the unique factorization

$$\mathrm{Spec}^{\mathcal{G}} R \to B^{\mathcal{G}}GL_j \to \coprod_{i\in I} B^{\mathcal{G}}GL_i.$$

Since the sheafification commutes with coproducts, $\coprod_{i\in I} B^{\mathcal{G}}GL_i$ is equivalent to the sheafification of the presheaf given by $R \mapsto \coprod_{i\in I}(B^{\mathcal{G}}GL_i(R))$. (Note that the coproduct of sheaves are the sheafification of the objectwise coproduct as presheaves, and an functor obtained by objectwise coproduct of sheaves is not always a sheaf.) Therefore, an object of $(\coprod_{i\in I} B^{\mathcal{G}}GL_i)(R)$ can be represented by a finitely generated projective R-module whose local ranks are contained in I.

Let $I\mathrm{Mod}_R^{proj}$ be an ∞-category of finitely generated projective modules whose local rank is contained in I. Since R is Zariski connected, by Lemma 4.23(iii), we deduce that $I\mathrm{Mod}_R^{proj} \simeq \coprod_{\{i\in I\}} \mathrm{Mod}_R^{iProj}$. Therefore, for any object of $(\coprod_{i\in I} B^{\mathcal{G}}GL_i)(R)$, there exists a unique $j \in I$ such that the object can be represented by a finitely generated projective R-module of rank j.

By applying the Yoneda embedding [8, Section 5.1.3], an object of $(\coprod_{i\in I} B^{\mathcal{G}}GL_i)(R)$ represented by a finitely generated projective R-module of rank j corresponds to the morphism $\mathrm{Spec}^{\mathcal{G}} R \to \coprod_{i\in I} B^{\mathcal{G}}GL_i$ which has the unique factorization

$$\mathrm{Spec}^{\mathcal{G}} R \to B^{\mathcal{G}}GL_j \to \coprod_{i\in I} B^{\mathcal{G}}GL_i.$$

□

We consider a decomposition of an $\mathbb{E}_\infty$-ring R by using the idempotent element in $\pi_0 R$.

Proposition 4.26. *Let* $\mathrm{Spec}^{\mathcal{G}} R : (\mathrm{CAlg}^{\mathcal{G}})^{op} \to \widehat{\mathcal{S}}$ *be a spectral scheme and* $\mathrm{Spec}^{\mathcal{G}} R \to B^{\mathcal{G}}GL_n$ *a morphism in* $\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})$. *Let* $B^{\mathcal{G}}GL$ *be a sheaf* $\coprod_{n\in\mathbb{N}} B^{\mathcal{G}}GL_n$. *For* $I \subset J \subset \mathbb{N}$, *we have the system* $\coprod_{i\in I} B^{\mathcal{G}}GL_i \to \coprod_{j\in J} B^{\mathcal{G}}GL_j$ *given by inclusions. Then there is an equivalence of* ∞*-groupoids;*

$$\mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\ B^{\mathcal{G}}GL) \simeq (\mathrm{Mod}_R^{\infty proj})^{\simeq}.$$

Proof. By Proposition 4.24, we have

$$\operatorname*{colim}_{I\subset\mathbb{N}} \mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\ \coprod_{i\in I} B^{\mathcal{G}}GL_i) \simeq \mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\ B^{\mathcal{G}}GL).$$

By decomposing R with the Zariski connected $\mathbb{E}_\infty$-rings (given by the corresponding irreducible decomposition on π_0) and applying Lemma 4.25, for $I = \{1, \cdots, n\}$, we have the equivalence

$$\operatorname*{colim}_{I\subset\mathbb{N}} \coprod_{i\in I} \mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\ B^{\mathcal{G}}GL_i) \simeq \operatorname*{colim}_{I\subset\mathbb{N}} \mathrm{Map}_{\mathcal{S}hv_{\widehat{\mathcal{S}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\ \coprod_{i\in I} B^{\mathcal{G}}GL_i).$$

By Proposition 4.21, we identify the left hand side with

$$\operatorname*{colim}_{i\in\mathbb{N}} (\mathrm{Mod}_R^{iproj})^{\simeq} \simeq (\mathrm{Mod}_R^{\infty Proj})^{\simeq}.$$

□

5. K-theory of $\mathrm{Mod}_R^{\infty proj}$

Definition 5.1 (cf. [2] 1.2, [4] Definition 2.1)**.** Let $\mathcal{C}$ be a pointed ∞-category. A class of w^∞-cofibrations is a class of morphisms in $\mathcal{C}$ which satisfies the following conditions:

(i) $* \to X$ is a w^∞-cofibration for any object X,

(ii) The class of w^∞-cofibrations includes weak equivalences,

(iii) Any composition of w^∞-cofibrations is a w^∞-cofibration,

(iv) For a w^∞-cofibration $X \to Y$ and a morphism $X \to Z$, there exists a pushout square

$$\begin{array}{ccc} X & \longrightarrow & Y \\ \downarrow & & \downarrow \\ Z & \xrightarrow[f]{} & W, \end{array}$$

in which the morphism f is a w^∞-cofibration.

We call such a pair of $\mathcal{C}$ and a w^∞-cofibrations an ∞-category with w^∞-cofibrations.

In terminology of [3], a w^∞-cofibration is called a cofibration. In terminology of [1], a w^∞-cofibration is called an igressive morphism.

Remark 5.2. We say that a w^∞-cofibration is a split w^∞-cofibration if it is split as a morphism.

Definition 5.3. We make into $\mathrm{Mod}_R^{\infty proj}$ to be an ∞-category with w^∞-cofibrations as follows.

Declare a morphism $f : P_1 \to P_2$ in $\mathrm{Mod}_R^{\infty proj}$ to be a w^∞-cofibration if it is a morphism in Mod_R and the cofiber of f is an object in $\mathrm{Mod}_R^{\infty proj}$.

Then, Mod_R^{proj} (resp. $\mathrm{Mod}_R^{\infty proj}$) is an ∞-category with w^∞-cofibrations.

Lemma 5.4. *(i) A w^∞-cofibration in $\mathrm{Mod}_R^{\infty proj}$ is always split, e.g., it is a split w^∞-cofibration in Remark 5.2.*

(ii) Let $R_1 \to R_2$ be a morphism of connective $\mathbb{E}_\infty$-rings. Then, a functor $\mathrm{Mod}_{R_1}^{\infty proj} \to \mathrm{Mod}_{R_2}^{\infty proj}$ given by the extension of coefficients is an exact functor in the sence of K-theory.

Proof. By [9, Proposition 7.2.2.6 (5)], a cofiber sequence of finitely generated projective modules is always split up to homotopy. Thus, (i) holds.

Since the tensor product commutes with the cofiber sequences and preserves rank, the extension of coefficients is exact. □

Remark 5.5. In this paper, we consider the connective K-theory. We can regard the K-theory spectrum as a K-theory space.

Let $\mathcal{C}$ be a pointed additive ∞-category. We recall an ∞-version of the additivity theorem [17, Theorem 1.8.7] given by [5]. We state the theorem as in Lurie's unpublished note [14].

Recall the notion of a split w^∞-cofibration from Remark 5.2 and the notion of the group completion. We will apply the following theorem for $\mathrm{Mod}_R^{\infty proj}$ whose w^∞-cofibrations defined in Definition 5.3 are split w^∞-cofibrations by Lemma 5.4.

Theorem 5.6 ([5], cf. [14] Theorem 10)**.** *Let $\mathcal{C}$ be a pointed additive ∞-category with split w^∞-cofibrations, and $\mathcal{C}^\simeq$ be the maximal ∞-groupoid of $\mathcal{C}$ as in (4.3). Let $K(\mathcal{C})$ be the algebraic K-theory and $\Omega B(\mathcal{C}^\simeq)$ the group completion of $\mathcal{C}^\simeq$.*

Then, there is an equivalence

$$\Omega B(\mathcal{C}^\simeq) \to K(\mathcal{C}) \tag{5.1}$$

of spases. Here, Ω and B are defined in Section 4.

Corollary 5.7. *We have $\Omega B((\mathrm{Mod}_R^{\infty proj})^\simeq) \simeq K(\mathrm{Mod}_R^{\infty proj})$.*

Proof. The ∞-category $\mathrm{Mod}_R^{\infty proj}$ (resp. Mod_R^{proj}) is additive, pointed by 0, with split w^∞-cofibrations. Thus, Theorem 5.6 can be applied. □

6. Proof of Theorem 1.2

Let us keep the notation explained in the previous sections. Now we prove Theorem 1.2.

Definition 6.1. Let $B^{\mathcal{G}}GL = \coprod_{n\in\mathbb{N}} B^{\mathcal{G}}GL_n$. We define a morphism $t : B^{\mathcal{G}}GL \times B^{\mathcal{G}}GL \to B^{\mathcal{G}}GL$ by the morphism induced from the assignment

$$B^{\mathcal{G}}GL_n \times B^{\mathcal{G}}GL_m \to B^{\mathcal{G}}GL_{n+m}$$

which sends a pair (P, Q) of projective modules of rank n and m respectively to the projective module $P \oplus Q$ of rank $n + m$ under the identification of Proposition 4.21.

Proposition 6.2. *Let $B^{\mathcal{G}}GL = \coprod_{n\in\mathbb{N}} B^{\mathcal{G}}GL_n$. Then, $B^{\mathcal{G}}GL$ becomes a commutative ∞-monoid by the morphism t defined in Definition 6.1 as the multiplication.*

Proof. For projective modules P and Q of rank n and m respectively, $P \oplus Q$ is the projective module of rank $n+m$. This assighment is obviously commutative and associative. □

Theorem 6.3. *Let $\mathrm{CAlg}^{\mathcal{G}}$ be the ∞-category CAlg^{cn} equipped with Zariski or Nisnevich topology defined in Section 2. Let $B^{\mathcal{G}}GL = \coprod_{n\in\mathbb{N}} B^{\mathcal{G}}GL_n$ and $\Omega B^{\mathcal{G}}(B^{\mathcal{G}}GL)$ the group completion as a sheaf on $\mathrm{CAlg}^{\mathcal{G}}$. Note that $B^{\mathcal{G}}GL$ becomes a commutative ∞-monoid as in Proposition 6.2.*

There is an equivalence of ∞-groupoids:

$$\mathrm{Map}_{Shv_{\widehat{\mathcal{G}}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\, \Omega B^{\mathcal{G}}(B^{\mathcal{G}}GL)) = \widetilde{K}^{\mathcal{G}}(\mathrm{Mod}_R^{\infty proj}),$$

where $\widetilde{K}$ is the sheafification of K defined by (1.1).

Proof. By Proposition 4.21, we have an equivalence

$$\Omega B(B^{\mathcal{G}}GL(R)) \simeq \Omega B((\mathrm{Mod}_R^{\infty proj})^\simeq),$$

where $(-)^{\simeq}$ denotes the maximal ∞-groupoid (4.3). Since all w^{∞}-cofibrations in $\mathrm{Mod}_R^{\infty proj}$ are split and the homotopy category of Mod_R^{proj} is additive, by Corollary 5.7, we obtain that $\Omega B((\mathrm{Mod}_R^{\infty proj})^{\simeq})$ is equivalent to the algebraic K-theory $K(\mathrm{Mod}_R^{\infty proj})$ given by $S_{\bullet}$ construction.

On the other hand, we have an equivalence induced by Yoneda embedding

$$\mathrm{Map}_{\mathcal{S}hv_{\mathcal{G}}(\mathrm{CAlg}^{\mathcal{G}})}(\mathrm{Spec}^{\mathcal{G}} R,\, \Omega B^{\mathcal{G}}(B^{\mathcal{G}}GL)) \simeq (\Omega B^{\mathcal{G}} B^{\mathcal{G}} GL)(R).$$

By virtue of Proposition 6.2, we can apply Proposition 3.15 to the commutative ∞-monoid sheaf $B^{\mathcal{G}}GL$. The sheafification of the objectwise group completion functor $R \mapsto \Omega B(B^{\mathcal{G}}GL(R))$ is equivalent to the group completion of the sheaf given by the assignment $R \mapsto (\Omega B^{\mathcal{G}} B^{\mathcal{G}} GL)(R)$. Thus, we have an equivalence $(\Omega B^{\mathcal{G}} B^{\mathcal{G}} GL)(R) \simeq \Omega B(B^{\mathcal{G}}GL(R))$ after the sheafification. □

References

[1] C. Barwick, *On the algebraic K-theory of higher categories*, Preprint, available at arXiv:1204.3607 (2014).

[2] ———, *On the exact ∞-categories and the theorem of the heart*, Preprint, available at arXiv:1212.5232v4 (2014).

[3] Andrew J. Blumberg, David Gepner, and Gonçalo Tabuada, *A universal characterization of higher algebraic K-theory*, Geom. Topol. **17** (2013), no. 2, 733–838, DOI 10.2140/gt.2013.17.733. MR3070515

[4] A. Blumberg, D. Gepner, and G. Tabuada, *The algebraic K-theory endomorphism*, available at arXiv:1302.1214 (2014).

[5] T. Fiore, W. Luck, and M. Pieper, *Waldhausen Additivity : Classical and Quasicategorical*, Preprint, available at arXiv:1207.6613 (2012).

[6] D. Gepner, M. Groth, and T. Nikolaus, *Universality of multiplicative infinite loop space machines*, Preprint, available at arXiv:1305.4550v1 (2013).

[7] D. Kodjabachev and S. Sagave, *Strictly commutative models for $\mathbb{E}_{\infty}$ quasi-categories*, Preprint, available at arXiv:1408.0116v2 (2014).

[8] J. Lurie, *Higher topos theory*, Annals of Mathematics studies, vol. 170, Princeton University Press, 2009.

[9] ———, *Higher algebra*, Preprint, available at www.math.harvard.edu/lurie (2011).

[10] ———, *Quasi-Coherent Sheaves and Tannaka Duality Theorems*, Preprint, available at www.math.harvard.edu/lurie (2011).

[11] ———, *Decsent theorems*, Preprint, available at www.math.harvard.edu/lurie (2011).

[12] ———, *Structured spaces*, Preprint, available at www.math.harvard.edu/lurie (2011).

[13] ———, *Spectral Schemes*, Preprint, available at www.math.harvard.edu/lurie (2011).

[14] ———, *Additive K-Theory (Lecture 18)*, lecture available at http://www.math.harvard.edu/∼lurie/281notes/Lecture18-Rings.pdf (2014).

[15] ———, *Algebraic K-Theory of Ring Spectra (Lecture 19)*, available at http://www.math.harvard.edu/ lurie/281notes/Lecture19-Rings.pdf (2014).

[16] F. Morel and V. Voevodsky, $\mathbf{A}^1$-*homotopy theory of schemes*, Institut des Hautes Études Scientifiques. Publications Mathématiques **90** (1999), 45–143 (2001).

[17] F. Waldhausen, *Algebraic K-theory of spaces*, Lecture Notes in Math. **1126** (1985), 318–419.

Department of Mathematical Sciences, Shinshu University, 3-1-1, Asahi, Matsumoto City, JAPAN, 390-8621.

E-mail address: primarydecomposition@gmail.com